



Effect of Sorghum-tigernut Ibyer (A Traditional Gruel) on the Fasting Blood Glucose Levels of Alloxan-induced Diabetic Rats

Shalem Shiekuma^{1*}, Moses Ukeyima¹, Msendoo, Janet Ahuah¹, Idoko Blessing¹ and Terzungwe Tughgba²

¹Department of Food Science and Technology, College of Food Technology and Human Ecology, Federal University of Agriculture, Makurdi, Benue State, Nigeria.

²Department of Theriogenology, College of Veterinary Medicine, Federal University of Agriculture, Makurdi, Benue State, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. Authors SS and MU designed the study, wrote the protocol and supervised the research. Author MJA managed the analyses of the study and performed the statistical analysis. Author IB managed the literature searches and wrote the first draft of the manuscript. Author TT managed all treatments and management of laboratory animals used for the study. All authors read and approved the final manuscripts.

Article Information

DOI: 10.9734/EJNFS/2019/v9i330065

Editor(s):

- (1) Dr. R. C. Ranveer Department of Meat, Poultry and Fish, PG Institute of Post – Harvest Management, India.
(2) Dr. Morten Poulsen, Head of Research Group, Div. of Toxicology and Risk Assessment, National Food Institute, Technical University of Denmark, Denmark.

Reviewers:

- (1) Mohini Chetan Kuchekar, Pune University, India.
(2) K. D. Mini, Mahatma Gandhi University, India.
(3) E. Siva Rami Reddy, Tanta University, India.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/49050>

Original Research Article

Received 13 April 2019
Accepted 19 June 2019
Published 29 June 2019

ABSTRACT

Background: There is growing interest in the use of natural foods in the management of chronic diseases like diabetes. Ibyer is a fibre rich gruel consumed amongst the Tiv people of Benue State made from whole sorghum or millet flours.

Aim: The aim of the study was to evaluate the effect of sorghum-tigernut ibyer on the fasting blood glucose levels and body weight of alloxan monohydrate-induced diabetic rats.

Methods: Sorghum flour (SF) and tigernut flour (TNF) were blended at different proportions

*Corresponding author: Email: shalemshie@gmail.com;

(100:00; 90:10; 80:20; 70:30) for the purpose of ibyer production. The flour samples were subjected to proximate analysis using standard analytical procedures, the sensory attributes of ibyer produced from the different flour samples was evaluated on a 9-point hedonic scale. Thirty (30) male Wistar rats (100–180 g body weight) were grouped into five (1-5) each group containing six rats. They were induced with diabetes by injecting them with 150ml/kg of body weight with alloxan monohydrate dissolved in saline water (0.9% NaCl) except for group 1. Blood samples were collected from the tail of the rats, prior to induction, 48hrs after induction and 72 hrs after three days of continuous feeding with test diet. Fasting blood glucose was measured using a standard glucometer and test strips.

Results: The sensory attributes indicated that ibyer produced from the flour samples were generally acceptable. Fasting blood glucose levels after 72 hrs of feeding were found to be lowered more in groups giving flours with a higher proportion of Tigernut.

Conclusion: The results indicated that sorghum-tigernut ibyer exerted hypoglycaemic effect on the experimental animals.

Keywords: Sorghum; tigernut; Ibyer; diabetic rats; blood glucose level.

1. INTRODUCTION

Diabetes is a chronic disease in which food (particularly the nutrient – carbohydrate) is not properly absorbed in the body to be used as energy [1]. Diabetes and hypertension are global health disorders afflicting millions of people worldwide with an ever-increasing incidence and prevalence the upsurge of diabetes in Africa has been linked with rapid urbanization and changing dietary habits [2,3]. Use of indigenous foods has been advocated to reduce the incidence of chronic, diet-related non-communicable diseases such as obesity, diabetes, cardiovascular diseases and stroke [4]. The development of diabetes mellitus, obesity, cancer and cardiovascular disease (CVD) has been reported to be linked to the intake of high glycaemic index (GI) foods while intake of low GI foods has been shown to play a positive role in the management of these diseases [5].

Ibyer is an indigenous non-alcoholic gruel made from cereals (maize, sorghum and millet), consumed in Nigeria especially in Benue state by Tiv people [6]. It is prepared by cooking reconstituted whole cereal flour or paste in water and is often served with beans products such as *akpukpa* (native bread) and beans cake. Sour type (*ibyer-i-angen*) arises as result of the fermentation step that is undertaken before the porridge is prepared. The flour or wet milled paste is usually reconstituted with water or form slurry or paste after which it is left for a given time to enable fermentative microbial organisms to act on the product, producing the characteristic sourness associated with the product. Traditionally, the fermentation time lasts overnight.

Sorghum (*Sorghum bicolor L. Moench*) is the fifth most important cereal after wheat, rice, maize and barley in terms of production and utilization. The total world annual sorghum production is over 60 million tons from cultivated area of 45 million for food, alcoholic beverages and the grain is one of the staple foods for poor and rural people. Sorghum is gluten-free thus can be consumed by people with celiac diseases [7]. The important nutrients of sorghum include; dietary fibre, fat-soluble and B-vitamins and minerals [8]. Sorghum flour is used for flours, porridge and side dishes, malted and distilled beverages and special food such as popped grain, its protein content is higher than many grains. Sorghum is rich in anti-oxidant which is believed to help lower the risk of cancer, diabetes, heart diseases and other neurological diseases.

Tigernut has been used extensively mainly for human consumption in Spain [9]. It was an important food in ancient Egypt [10]. Nowadays tigernuts are cultivated in Northern Nigeria, Mali, Senegal, Ahana and Togo where they are used primarily uncooked as a side dish [11]. The flour is a good alternative for other flour like wheat flour, as it is gluten free and good for people who cannot take gluten in their diets. It is also used to make cakes and biscuits and the oil is used for cooking [12]. The dietary fibre content of tigernut is effective in the treatment and prevention of diseases such as colon cancer, coronary heart diseases, obesity, diabetes and gastro-intestinal disorders [13]. This research was aimed at determining the effect of sorghum-tigernut '*Ibyer*' on fasting blood glucose levels of alloxan monohydrate induced diabetic rats.

2. MATERIALS AND METHODS

2.1 Sample Preparation

The red sorghum grain variety and dried tigernuts were purchased from Wadata Market in Makurdi, Benue State, Nigeria. Alloxan monohydrate was purchased from Sigma Chemical Co. (St. Louis, MO, USA).

2.2 Sample Preparation

2.2.1 Sorghum flour

The preparation of sorghum flour was done according to the method described by Sergio [14]. Sorghum grains were dry-cleaned by handpicking, washed in a large volume of water to remove impurities and sun-dried. The dried sorghum grains were milled into sorghum flour using hammer mill.

2.2.2 Tigernut flour

Tigernut flour was prepared according to the method described by Adejuyitan et al. [15]. Dry tigernuts (brown variety) were sorted to remove unwanted materials like stones, pebbles and other foreign seeds before washing with tap water. The cleaned nuts were sun dried to a moisture content of about 13% then milled and sieved.

2.3 Formulation of Flour Blends

The two flours (SF and TNF) were blended in a ratio of 100: 0, 90:10, 80:20 and 70:30 respectively.

2.4 Preparation of *Ibyer-i-angen*

After the preparation of the sorghum-tigernut flour blends, the flour samples were mixed with 200 ml of water in five different containers with a cover and kept at ambient temperature ($30\pm 2^{\circ}\text{C}$) for 24 hrs to allow natural fermentation take place. The fermented paste from the two flour blends was diluted with water and cooked in a tower aluminium pot containing about 80 cl of water on a gas cooker. The mixture was stirred for about 7-8 min to achieve desired consistency. For feeding to the animals, the prepared/cooked *ibyer* was oven dried and milled.

Flow chat for production of *ibyer-i-angen* is shown in Fig.1

2.5 Determination of Proximate Composition

The samples were separately analysed for moisture, ash, fat, and fibre contents respectively using standard methods of the Association of Official Analytical Chemists [16]. Crude protein contents ($\text{N} \times 6.25$) were estimated from the crude nitrogen contents of the samples, as determined using Micro Kjeldahl method [17]. Carbohydrate content of the sample was determined by difference as described by Ihekoronye [18].




Fig. 1. Traditional method of production of *Ibyer-i-angen*

2.6 Sensory Analysis

Sensory characteristics of *ibyer-i-angen* prepared from sorghum and tigernut was evaluated and assessed by 20 semi-trained panellists of Department of Food Science and Technology, University of Agriculture, Makurdi who are familiar with *ibyer-i-angen*. Fresh samples of cooked porridge/gruel were assessed for appearance, taste, mouth feel, flavour and general acceptability. The judges recorded the quality characteristics of each sample using nine-point hedonic scale, where 1 = dislike extremely, 5 = neither like nor dislike, and 9 = like extremely as described by Iwe [19].

2.7 Animal Treatments

Thirty (30) male Wistar rats (100–180 g body weight) were obtained from the small Animal Experimental Unit of the National Veterinary Research Institute Vom, Plateau State. The rats were housed under standard hygienic conditions in metal cages with wood shavings as bedding. Rats were also kept under natural thermal environmental conditions with ambient

temperature of 24 C–26 C and relative humidity of 70%–80%, and approximately alternating 12 hr light/dark cycles. They were given access to a standard pelletized rat chow and water *ad libitum*. The animal experiments adhered to the Guide for the Care and Use of Laboratory Animals.

2.8 Induction of Experimental Diabetes

Diabetes mellitus (DM) was induced in overnight-fasted rats by a single i.p. injection of freshly-prepared alloxan monohydrate, dissolved in a cold physiological saline (0.9% NaCl) solution at the dose rate of 150 mg/kg body weight. The animals were given free access to 5% glucose solution in order to overcome the alloxan-induced hypoglycaemia for the first one-hour post-treatment with alloxan monohydrate. Blood glucose concentration of the rats was estimated 48 hours after alloxan administration and DM was confirmed by analysis of blood samples, collected from the vein at the tip of the tail, using a portable blood glucometer and glucose test strips (On Call®Plus, Hannover, Germany). Animals with blood glucose concentration equal or more than 14 mmol/L were considered diabetic and used in the entire experimental group. Animals were weighed and randomly assigned to six groups and treated as follows:

Group 1, Control group: (normal saline only);

Group 2, Diabetic + Treated with Sample A (alloxan 150 mg/kg, i.p);

Group 3, Diabetic + Treated with Sample B (alloxan 150 mg/kg, i.p);

Group 4, Diabetic + Sample C (alloxan 150 mg/kg, i.p);

Group 5, Diabetic + Sample D (alloxan 150 mg/kg, i.p);

Experimental animals were fed thirty grams (30 g) each of the test diet, after 48hrs of verification of diabetes, daily for 3 days.

2.9 Measurement of Blood Glucose

Experimental animals were rearranged according to the blood glucose concentration, except the control group, before commencement of treatment. Blood glucose concentration in all experimental groups were

recorded following overnight fasting 72 h after commencing the feed trials, using a portable glucometer (On Call®Plus, Hannover, Germany) and glucose test strips.

2.10 Measurements of Body Weight

Rats were weighed individually at Day 0 (before induction of DM), day 2 (post induction of DM) and 72 h (post-treatment with feed samples) using a digital precision weighing balance, and the body weights were recorded to calculate the body weight gains.

2.11 Statistical Analysis

Data collected were subjected to Analysis of Variance (ANOVA). Means were separated with Fisher's LSD using SPSS software (2009 model) and judged significantly different at 95% confidence level ($p < 0.05$).

3. RESULTS AND DISCUSSION

3.1 Results

3.1.1 Proximate composition of blends of sorghum and tigernut flours

Table 1 shows the results obtained for the proximate composition of blends of sorghum and tigernut flours. Moisture, protein and carbohydrate showed a decreasing trend with increasing incorporation levels of Tigernut flour. For the rest of the parameters the trend was opposite.

3.1.2 Sensory properties of lbyer produced from blends of sorghum and tigernut flours

The results of the sensory evaluation of *lbyer* produced from blends of sorghum and tigernut flours are shown in Table 2. The sensory qualities measured were appearance, mouth feel, taste, flavour and general acceptability. All the samples were generally acceptable though sample B was most preferred.

3.1.3 Effect of sorghum-tigernut lbyer on the fasting blood glucose levels of alloxan induced diabetic rats

The results of the fasting blood glucose levels of alloxan induced diabetic rats before and after feeding with test diets are presented in Fig. 2. The results showed that the average fasting glucose levels of the diabetic rats before feeding

with the test diets ranged from 22.30 mmol/L to 32.00 mmol/L and 3.15 mmol/L to 6.05 mmol/L in the non-diabetic rats (control group). However, there was considerable reduction in average fasting blood glucose levels of the diabetic rats 0hrs through to 72 hrs post treatment. The results ranged from 5.52 mmol/L from group 1 (control group) through to 5.45 mmol/L in group 5 with rats'

group 2 having the highest average fasting blood glucose levels of 17.60 mmol/L.

3.1.4 Body weight of rats before and after feeding with test diet

Fig. 3 presents results of body weight changes during the experimental period. Results showed

Table 1. Proximate composition of blends of sorghum and tigernut flours

Parameters (%)	Samples				LSD
	A	B	C	D	
Moisture	10.1 ^a ±0.02	8.2±0.01 ^b	6.2 ^c ±0.01	5.5 ^d ±0.01	0.02
Protein	7.5 ^a ±0.00	7.2 ^b ±0.00	7.0 ^c ±0.00	6.3 ^d ±0.01	0.02
Fat	3.2 ^e ±0.01	4.8 ^d ±0.00	8.4 ^c ±0.00	10.2 ^b ±0.00	0.01
Fibre	1.0 ^d ±0.03	2.1 ^c ±0.03	2.7 ^b ±0.01	3.2 ^a ±0.02	0.04
Ash	2.1 ^d ±0.01	2.4 ^c ±0.00	2.7 ^b ±0.01	2.8 ^a ±0.00	0.01
Carbohydrate	76.1 ^a ±0.02	75.3 ^b ±0.01	73.0 ^c ±0.02	72.1 ^d ±0.01	0.03

Mean with different superscript within the same column are significantly different at (P<0.05).
 LSD: Least Significant Difference SF: Sorghum Flour; TNF: Tigernut flour.
 A: 100% SF. B: 90% SF and 10% TNF C: 80% SF and 20% TNF D: 70% SF and 30% TNF

Table 2. Results of the sensory scores of "lbyer" produced from blends of sorghum and tigernuts flour

Sample	A	B	C	D	LSD
Appearance	7.15 ^a	7.05 ^a	6.85 ^a	6.20 ^{ab}	0.94
Flavour	6.65 ^a	6.75 ^a	6.75 ^a	6.05 ^a	1.17
Mouth feel	7.15 ^a	6.95 ^a	6.30 ^{ab}	5.60 ^{bc}	0.96
Taste	6.90 ^a	6.75 ^a	6.65 ^a	6.50 ^a	1.17
General acceptability	7.40 ^a	7.15 ^a	6.75 ^a	6.45 ^a	1.02

Mean with different superscript within the same column are significantly different at (P<0.05).
 LSD: Least Significant Difference; SF: Sorghum Flour; TNF: Tigernut flour.
 A: 100% SF. B: 90% SF and 10% TNF C: 80% SF and 20% TNF D: 70% SF and 30% TNF




Fig. 2. Fasting blood glucose levels of alloxan induced diabetic rats before and after feeding with test diet

Group 1(control; normal saline) Group 2(sample A at 100%SF) Group 3(sample B at 90%SF and 10%TNF)
 Group 4(sample C at 80%SF and 20% TNF) and Group 5 (sample D at 70%SF and 30%TNF)
 Key:SF; Sorghum flour; TNF: Tigernut flour




Fig. 3. The average body weight of the rats (grams)

Group 1(control; normal saline) Group 2(sample A at 100%SF) Group 3(sample A at 90%SF and 10%TNF)
Group 4(sample C at 80%SF and 20% TNF) and Group 5 (sample D at 70%SF and 30%TNF)

Key: SF; Sorghum flour; TNF: Tigernut flour

a weight increase in the non-diabetic rat group from 98 to 118 g from 0hrs to 72 hrs post feeding. However, the diabetic rat groups showed slight variations in body weight changes.

3.2 Discussion

3.2.1 Proximate composition of sorghum and tigernut flour blends

The moisture content of the sample flour decreased with increase in the tigernut flour substitution, which is due to the low moisture content found in tigernuts. The result is in agreement with those of [20] who reported a decrease in moisture content in flour. Low moisture content is an indication of good keeping quality of any food product; it also reduces microbial growth thereby extending the shelf life of the product.

The protein content decreased with increase in tigernut flour, which is attributed to the low protein content of tigernut as reported by Ade-Omomowaye et al. [21]. The protein content of the four samples meets the recommended dietary allowance (RDA) of 0.8g of protein per kilogram of body weight per day [22].

The fat content of the samples increased steadily from samples A to D with increase in tigernut substitution, as tigernut has been reported to be a rich source of fats [21,23,24,25]. Sorghum flour

has been reported to contain fats of up to 4.25%, [18,26]. The increase in the fat content could be attributed to the increase in the tigernut substitution. The fibre content of samples also increased with increase in tigernut substitution. These results were in accordance with [21] who reported an increase in the fibre content of tigernut-wheat flour with increasing tigernut flour substitution. There was a significant ($P < .05$) difference in the ash content of all the samples as the level of tigernut substitution increased. The ash content result is in agreement with the reports of [27] and comparable with those of [28]. Ash content is an indication of the mineral content in flour.

There was however, a decrease in the Carbohydrate content of Samples with a corresponding increase in tigernut substitution. There was a significant ($P < .05$) difference in carbohydrate content of all the samples. The decrease in carbohydrate content is attributed to an increase in tigernut flour.

3.2.2 Quality characteristics on the sensory attributes of lbyer produced from blends of sorghum and tigernut flour

The significant ($P < .05$) difference in sample with the highest proportion of Tigernut could be attributed to the fact that tigernut has a colour that is distinct from that of sorghum. It was however noted that the sensory scores for

appearance were considerable high even though with slight differences. This means all products were appealing to the eyes.

The significant difference in mouth feel in sample C and D could be attributed to higher amounts of fibre in samples C and D. *Ibyer* is a gruel made from whole cereals, particularly sorghum and millet, due to this it has a coarse feel when consumed.

There was also no significant ($P < .05$) difference among the samples in terms of taste, flavour and general acceptability. All the samples were generally acceptable indicating that *ibyer* of acceptable eating qualities can be produced from flour blends and sorghum and tigernuts. These results are in agreement with the reports of [6] who reported general acceptability of *ibyer* produced from sorghum and soy addition meaning addition of other legumes to the product may not affect the eating qualities of the product.

3.2.3 Effect of sorghum-tigernut lbyer on the fasting blood glucose levels of alloxan induced diabetic rats

The average fasting blood glucose of the rats in the various groups differed significantly. After three days of consuming approximately 30 g of the test diets daily, their fasting blood glucose levels dropped significantly, showing that rat groups fed with samples containing higher amounts of tigernut had lower fasting blood glucose levels than those of the control group. This could be attributed to the fact that tigernut is a rich source of fibre [29,30]. Fibre has been reported to exert some hypoglycaemic effects in subjects with type II diabetes. It is also reported that arginine an amino acid (not determined) found in tigernut has the potential of stimulating the release of insulin thereby ameliorating the effect of diabetes [31]. This could be one of the reasons for the lowered blood glucose levels after three days of consecutive feeding with the test diet. Several studies have shown that sorghum extracts and sorghum rich diets exert hypoglycaemic effects in either human subjects or lab animals, thus agreeing with the findings of this research [32,33].

4. CONCLUSION

Results of the sensory evaluation of *ibyer* produced from sorghum and tiger flour blends showed that all the samples were generally acceptable. The test diet was observed

to have some hypoglycaemic effect on the experimental animals, however, mechanism of action of achieving these results was not studied and so remains unclear. Therefore, it would be worthwhile, to further investigate the effect of this diet rich in Tigernut on blood glucose levels.

From the results, it is evident that nutritious diets can be formulated by complementing unexploited tuber and cereals like tigernut and sorghum respectively, such blends could be used to diversify their uses to develop new products.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Bellows L, Nichols K. Position statement of the American Diabetes Association. Nutrition Recommendation and intervention for diabetes: Diabetes Care. 2012;31:1.
2. Animaw W, Seyoum Y. Increasing prevalence of diabetes mellitus in a developing country and its related factors. PLoS One [Internet]. 2017;12(11): e0187670–e0187670. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29112962>
3. WHO. WHO | Double burden of malnutrition: Infographics. WHO [Internet]; 2017. [Cited 2019 Feb 7] Available: <https://www.who.int/nutrition/double-burden-malnutrition/en/>
4. Checkley W, Ghannem H, Irazola V, Kimaiyo S, Levitt NS, Miranda JJ, et al. Management of NCD in low- and middle-income countries. Glob Heart [Internet]. 2014;9(4):431–43. Available: <https://www.ncbi.nlm.nih.gov/pubmed/25592798>
5. McCulloch DK, Nathan DM, Lipman TO. Nutritional considerations in type 2 diabetes mellitus. 2016;1–26. [Cited 2019 May 7] Available: <https://www.uptodate.com/contents/nutritional-considerations-in-type-2-diabetes-mellitus>
6. Kure OA, Wyasu G. Influence of natural fermentation, malt addition and soya-fortification on the sensory and physico-

- chemical characteristics of Ibyer-Sorghum gruel. *Adv Appl Sci Res* [Internet]. 2013; 04(1):345–9
[Cited 2019 May 7]
Available:<http://www.imedpub.com/abstract/influence-of-natural-fermentation-malt-addition-and-soyafortification-on-thernsensory-and-physicochemical-characteristics-of-ibyersorghum-gruel-14775.html>
7. Taylor JRN, Achoch TJ, Bean SR. Novel food and non-food uses for sorghum and millets. *Articles in press. Journal of Cereal Science*. 2006;44:252-271.
 8. Waniska RO, Poe JH, Bandy OR. Effects of growth conditions on grain molding and phenols in sorghum *Caryopsis*. *Journal of Cereal Science*. 2004;10:217-225.
 9. Gambo A. Sensory evaluation of yoghurt produced from blend of Cowmilk, Soymilk and Tigernut milk. In: *Proceedings of 36th Annual Conference/AGM of Nigerian Institute of Food Science and Technology, Lagos*. 2012;440- 441.
 10. Neigbi OA. Quality characteristics of Candies Produced from Tigernuts Tubers (*Cyperus esculentus*) and Melon Seeds (*Citrulus colocynthis*) Milk Blends. *Global Journal of Science Frontier Research: Agriculture and veterinary*. 1992;15(2).
 11. Omode MO, Ibiom OF, Okoi A. Studies on the fungi and phytochemical and proximate composition of dry and fresh tigernuts (*Cyperus esculentus*. L). *International Research Journal of Biotechnology*. 2009;4(1):11-14.
 12. Adebayo-Oyetero AO, Ogundipe OO, Lofinmakin FK, Akinwande FF, Aina DO, Adeyeye SAO. Production and acceptability of chinchin snack made from wheat and tigernut (*Cyperus esculentus*) flour. *Cogent Food Agric* [Internet]. 2017;3(1):1282185.
Available:<https://www.tandfonline.com/doi/abs/10.1080/23311932.2017.1282185>
 13. Forouhi NG, Misra A, Mohan V, Taylor R, Yancy W. Dietary and nutritional approaches for prevention and management of type 2 diabetes. *BMJ* [Internet]. 201813;361:k2234.
Available:<http://www.bmj.com/content/361/bmj.k2234.abstract>
 14. Sergio Serna-Saldivar. *Cereal grains: Properties, processing and nutritional attributes* CRC Press; 2010. [ISBN 97814398156012010]
 15. Adejuyitan JA, Otunolam ET, Akandem ET, Bolarinwa IF and Oladokun FM. Some physicochemical properties of flour obtained from fermentation of tigernut (*Cyperus esculentus*) sourced from a market u Ogbomoso, Nigeria. *African Journal of Food Science*.2009;3(2):051-055.
 16. AOAC. *Official methods of Analysis, Association of Official Analytical Chemist* Washington, D. C; 2010.
 17. AOAC. *Official methods of Analysis, Association of Official Analytical Chemist* Washington, D. C; 2005.
 18. Ihekoronye, Ngoddy. *Integrated Food Science and Technology for the Tropics*; 1985.
 19. Iwe MO. *Hand book of sensory evaluation methods and analysis* Rojoint communication services Ltd. Enugu. 2002; 72-75.
 20. Abiose SH, Ikujenlola AV. Comparison of chemical composition, functional properties and amino acids composition of quality protein maize and common maize (*Zea may L.*); 2014.
 21. Ade-Omomowaye BIO, Akinwande BA, Bolarinwa IF, Adebisi AO. Evaluation of tigernut (*Cyperus esculentus*) - wheat composite flour and bread. *African Journal of Food Science*. 2008;(2):087-091.
 22. Yemi B. *Malnutriton; The case of Death In Nutrition*. 2010;785.
 23. Estoshola E, Oraedu AC. Fatty acid composition of Tigernut tubers, Baobab seeds (*Adansonia digitata* L.) and their mixtures. *American Journal of Chemical Sciences*. 2012;73:255-267.
 24. Zahras A, Ahmed MS. Explaining the suitability of incorporating tigernut flour as novel ingredient in gluten-free biscuit. *Journal of Food and Nutritional Sciences*. 2014;64(1):27-36.
 25. Nwaignikpe R. The phytochemical proximate and amino acid composition of the extracts of two varieties of tigernut (*Cyperus esculentus*) and their effects on sickle cell hemoglobin polymerization. *J. Med. Med. Sci*. 2010;1:543-549.
 26. Ibrahim FS, Babiker EE, Yousif NE, El-Tinay AH. Effect of whey protein supplementation and/or fermentation on biochemical and sensory characteristics of sorghum flour. *J. Food Technol*. 2005;3: 118-125.
 27. Adebowale AA, Adegoke MT, Sanni SA, Adegunwa MO, Fetuga GO. *Functional*

- properties and biscuit making potentials of sorghum - wheat Flour Composite. American Journal of Food Technology. 2012;7:372-379.
28. Aremu MO, Bamidele TO, Agere H, Ibrahim H, Aremu SO. Proximate composition and amino acid profile of raw and cooked black variety of tigernut (*Cyperus esculentus* L.) grown in Northern Nigeria. Journal of Biology Agriculture and Health Care. 2015;5:7.
29. Sánchez ZE, Fernández LJ, Angel Pérez AJ. Tigernut (*Cyperus esculentus*) commercialization: Health aspects, composition, properties and food applications. Comprehensive Reviews in Food Science and Food Safety. 2012;11(4):366–377.
30. Gambo A, Da' (). Tigernut (*Cyperus esculentus*): composition, products, uses and health benefits – a review. *Bayero journal of pure and applied sciences*. Vol. 2014;7(71):56–61.
31. Wu GY, Meininger CJ. Arginine nutrition and cardiovascular function. Journal of Nutrition. 2000;130(11):2626–2629.
32. Chung IM, Kim EH, Yeo MA, Kim SJ, Seo M and Moon HI. Antidiabetic effects of three Korean sorghum phenolic extracts in normal and streptozotocin-induced diabetic rats. Food Research International. 2011;44(1):127–132.
33. Park JH, Lee SH, Chung IM, Park Y. Sorghum extract exerts an anti-diabetic effect by improving insulin sensitivity via PPAR-?? in mice fed a high-fat diet. Nutrition Research and Practice. 2012;6(4):322–327. Available:https://doi.org/10.4162/nrp.2012.6.4.322

© 2019 Shiekuma et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://www.sdiarticle3.com/review-history/49050>